arXiv:nucl-th/0111033v1 12 Nov 2001

Chiral NN model and A_y puzzle

D. R. Entem* and R. Machleidt†
Department of Physics, University of Idaho, Moscow, Idaho 83844

H. Witała‡
Institute of Physics, Jagiellonian University, PL-30059 Cracow, Poland

(February 8, 2008)

We analyze the results by chiral NN models for the two-nucleon system and calculate the predictions for the nucleon vector analyzing power of elastic nucleon-deuteron (Nd) scattering, A_y, by these models. Our conclusion is that a *quantitative* chiral two-nucleon potential does not resolve the Nd A_y puzzle (when only two-body forces are included).

PACS numbers: 21.30.+y, 21.45.+v, 25.10.+s, 27.10+h

The term A_y *puzzle* refers to the inability to explain the nucleon vector analyzing power A_y in elastic nucleon-deuteron (Nd) scattering below 30 MeV laboratory energy for the incident nucleon by means of three-body calculations in which only two-nucleon forces are applied. The problem showed up as soon as it was possible to conduct three-body continuum calculations *with realistic NN potentials*. The first such calculation was performed by Stolk and Tjon [1] in 1978 using the Reid soft-core potential [2], and the first calculations with (a separable representation of) the Paris potential [3] were conducted by the Graz-Osaka group in 1987 [4]; in both cases, the A_y predictions showed the characteristic problem. Finally, the 'puzzle' became proverbial when rigorous three-nucleon continuum Faddeev calculations using realistic forces were started on a large scale [5]. Over the years, many measurements and calculations of Nd A_y were performed (including the pd reaction that involves the Coulomb force [6]) which all confirmed that the problem was real (see Ref. [7] for a review): For energies below 20 MeV, the A_y is predicted about 30% too small in the angular region around 120 deg center-of-mass angle where the maximum occurs.

There have been many attempts to solve the problem. Already in the very early stages of three-body continuum calculations, when only schematic NN potentials were applied, it was noticed that the Nd A_y predictions depend very sensitively on the strength of the input NN potential in the triplet P waves [8,9]—a sensitivity that was confirmed in later calculations using realistic forces [10]. Based upon this experience, Witała and Glöckle [11] showed in 1991 that small changes in those ^3P wave potentials could remove the discrepancy. This finding gave rise to systematic investigations of the question whether the small variations of the low-energy phase shifts of, particularly, those triplet P waves necessary to explain the Nd A_y are consistent with the NN data base. While Tornow and coworkers [12] suggest that the low-energy NN data may leave some lattitude in the NN ^3P waves that could

*On leave from University of Salamanca, Spain. Electronic address: dentem@uidaho.edu

†Electronic address: machleid@uidaho.edu

‡Electronic address: witala@if.uj.edu.pl

improve the predictions for Nd A_y, Hüber and Friar [13] find that it is not possible with reasonable changes in the NN potential to increase the Nd A_y and at the same time to keep the two-body observables unchanged.

Another important observation has been that conventional three-nucleon forces (when added to a realistic two-nucleon potential) change the predictions for Nd A_y only slightly and do not improve them [14,6]. Therefore, the general perception in the community has shifted towards the believe that the A_y puzzle is the 'smoking gun' for new types of three-nucleon forces [15–18] or new physics [19].

However, very recently, there has been an apparent indication that the above conclusion/believe may be premature. It was reported [20] that with a two-nucleon force of a new type, namely, one that is based upon chiral effective field theory, the A_y puzzle is resolved.

In recent years, effective field theory methods have become increasingly popular in nuclear physics. The reason for this development is the need to link conventional nuclear physics methods one way or the other to the underlying theory of strong interactions, QCD. After quark cluster models had only a limited success, it was recognized that the symmetries of QCD are more important than the high-energy degrees of freedom of QCD (quarks and gluons). The effective field theory concept distinguishes between different energy scales and assignes appropriate degrees of freedom for each scale while observing the over-all symmetries. For traditional nuclear physics with energies below 1 GeV, the right degrees of freedom are nucleons and pions interacting via a force that is controlled by (broken) chiral symmetry.

The derivation of the nuclear force from chiral effective field theory was initiated by Weinberg [21] and pioneered by Ordóñez [22] and van Kolck [23,24]. Subsequently, many researchers became interested in the subject [25–36]. As a result, efficient methods for deriving the nuclear force from chiral Lagrangians emerged and the quantitative nature of the chiral NN potential improved. This trend shows up, in particular, in the excellent work by Epelbaum *et al.* [30] where the chiral NN force was constructed using a unitary transformation and applying systematic power counting in next-to-leading order (NLO) and NNLO; and it is this potential in NLO that was applied in Ref. [20], resulting—seemingly—in a resolution of the long-standing A_y puzzle. It is the purpose of this note to critically investigate the predictions by the chiral NN model and the implications for the Nd A_y puzzle.

We start our investigation by taking a close look at important phase shifts of two-nucleon scattering. In Table I, we list S-wave phase shifts and, in Table II, we show ^3P-wave phase shifts for energies between 1 and 200 MeV. Since charge-dependence of the NN interaction is not a crucial factor in the A_y puzzle [12], and since the present chiral NN potentials are all charge-independent and adjusted to the neutron-proton (np) data, we consider np phase shifts and np data.

It is of interest to compare the phase shifts produced by the chiral NLO (next-to-leading order) model by Epelbaum *et al.* [30] (which is the chiral potential applied in Ref. [20]) with the empirical ones from the Nijmegen multi-energy np analysis [37] (PWA93) and the predictions by one representative of the family of the high-precision potentials constructed in the 1990's (CD-Bonn [38,39]). In S waves (Table I), there is, generally, good agreement up to 50 MeV. Above 50 MeV, differences between NLO and PWA93 show up and increase with energy. However, since the S-waves are not very important for Nd A_y, this may not have much impact on the predictions.

We turn now to the triplet-P waves (Table II) which are crucial for Nd A_y, and focus, first, on the energy range below 30 MeV. The NLO $^3P_0$ phase shifts are about 2% larger than PWA93, which is not significant. For $^3P_1$, the differences are more drastic: The NLO value at 10 MeV is about 5% smaller and the one at 25 MeV is 3% smaller than PWA93. Finally, the NLO prediction for $^3P_2$ at 25 MeV is enhanced by 13%.

To understand how variations of the ^3P phase shifts may effect observables, we consider the spin-orbit phase shift combination,

$$\Delta_{\rm LS} = \frac{1}{12}\left(-2\,\delta_{^3P_0} - 3\,\delta_{^3P_1} + 5\,\delta_{^3P_2}\right)\,, \tag{1}$$

which is a measure for the strength of the spin-orbit force. Results are shown in Table III. At 10 MeV, one obtains $\Delta_{\rm LS}^{\rm PWA93} = 0.203$ deg and $\Delta_{\rm LS}^{\rm NLO} = 0.212$ deg for PWA93 and NLO, respectively, implying that the NLO value is larger by 4.4% as compared to PWA93. At 25 MeV, the corresponding figures are $\Delta_{\rm LS}^{\rm PWA93} = 0.93$ deg and $\Delta_{\rm LS}^{\rm NLO} = 1.09$ deg implying that NLO is 17% larger. These numbers show that the spin-orbit force of NLO is enhanced as compared to the Nijmegen analysis and similar enhancements are obtained when comparing to any of the high-precision potentials, like CD-Bonn (cf. Table III). It is well-known that the two-nucleon spin-orbit force is magnified in the Nd system. Therefore, a moderate enhancement of the NN LS-potential leads to a substantial enlargement of the Nd A_y. Thus, the enhanced NLO spin-orbit force as reflected in the larger values for $\Delta_{\rm LS}^{\rm NLO}$ could very well be the explanation of the large Nd A_y predictions by NLO reported in Ref. [20].

Next, we consider the ^3P phase shifts above 30 MeV. Here, the differences between PWA93 and CD-Bonn, on the one hand, and NLO, on the other, are in general larger and increase with energy (cf. Table II). This trend is most dramatic in $^3P_2$ where the discrepancies quickly grow into the hundreds of standard deviations. The sign of these differences are such as to drastically enhance the spin-orbit force, see Table III for the value at 50 MeV. Note that the NN t-matrix, on- and off-shell, is input to the three-body continuum calculations. Thus, the description of the two-nucleon data at energies above 30 MeV has impact on low-energy three-body predictions. Therefore, the unrealistically strong NLO spin-orbit force above 30 MeV may—by means of an off-shell effect—further enhance the Nd A_y predictions. However, one may not have much confidence in this type of off-shell effect.

The above observations trigger the question if chiral models can also make more accurate predictions for NN phase shifts; and if so, what are the implications for the Nd A_y problem, in such a case. The natural way is to include higher order terms in power counting which should improve not only the quality of the NN phase shift reproduction but also extend the energy range in which it works. For that purpose, we pick up the chiral NN potential of Ref. [40] (subsequently denoted by 'Idaho') that was recently developed. In the chiral Idaho model [40], contact terms up to order four are included which introduces more parameters allowing for a better fit of the lower partial waves in a much wider energy range. In Table I and II, it is clearly seen that the chiral Idaho NN potential reproduces the empirical np phase shifts of PWA93 up to 200 MeV, accurately.

We consider now the observable Nd A_y which is the focus of this paper. We have calculated the predictions for Nd A_y at energies 3, 10, and 65 MeV for the incident nucleon. The results are shown in Fig. 1. In this figure, the shaded band represents the prediction by the family of high-precision potentials (using always the np version of those models), namely, CD-Bonn [39], Argonne V_{18} [42], and the Nijmegen potentials Nijm-I, Nijm-II, and

Nijm93 [41]. The dashed line is the predicition by the Idaho chiral NN potential [40] and it is clearly seen that this prediction follows accurately the narrow band made up from the variations among those high-precision potentials. In conclusion, at 3 and 10 MeV, we are still having an A_y problem if the chiral NN potential is a *quantitative* one.

The evidence presented may be perceived as a convincing proof that a *quantitative* chiral potential does not resolve the Nd A_y puzzle. However, there remains one objection that can be raised. In the literature, notably in Ref. [12], one can find the suggestion that the ^3P waves at low energy are not as well determined as claimed in PWA93 [37]. If true, then moderate variations of the ^3P phase shifts at low energy could be consistent with the low-energy NN data. This variations could be such as to enhance the low-energy spin-orbit force and, thus, lead to an improved prediction for Nd A_y. For the purpose of seriously checking out this possibility, we have constructed a variation of the Idaho chiral potential with modified ^3P phase shifts at low energy. The column 'Modified' of Table II shows the ^3P phase shifts and the corresponding column of Table III reveals that for this fit the spin-orbit force is enhanced, similarly to NLO. However, in contrast to NLO, the 'modified' model is much more realistic, since the phase shifts do not diverge to unrealistic values at higher energies.

We have calculated the Nd A_y as predicted by the 'modified' chiral model and find, indeed, a considerable improvement (see dotted curve in Fig. 1). Is this the resolution of the A_y puzzle by a chiral NN potential? To answer this question one needs to know if the modified chiral model is a realistic and quantitative NN potential. A precise and reliable answer cannot be given by just looking at phase shifts. As stressed repeatedly by the Nijmegen group in the past, only a direct confrontation with the NN data can reveal if a NN potential is quantitative or not. For that reason we have calculated the χ^2/datum for the fit of the world np data as represented by the 1999 database [43], see Table IV. For a proper interpretation of the results of Table IV, it is necessary to establish a standard concerning what χ^2/datum represents a 'quantitative' reproduction of the data. This issue was debated a lot in the 1990's, and the consensus that emerged was that only values below 1.1 are acceptable. Deliberatly, we losen this standard and consider a fit with $\chi^2/\text{datum} \leq 1.2$ as 'quantitative', while we will perceive higher values as not acceptable.

Applying this standard, the modified model produces unacceptable values for χ^2/datum for all energy intervals below 75 MeV (cf. Table IV). Thus, the modified chiral model is *not* a quantitative one and, consequently, it is *not* the resolution of the Nd A_y puzzle. Since it is well known that the off-shell character of the NN potential plays essentially no role in three-nucleon scattering, one can further draw the more general conclusion: No model that reproduces the NN data correctly can solve that Nd A_y puzzle.

Table IV shows also the χ^2/datum of the other models discussed in this paper. It is seen that PWA93, Idaho, and CD-Bonn reproduce the np data below 210 MeV with the perfect $\chi^2/\text{datum} = 0.97$ and 0.98. The chiral NLO potential by Epelbaum *et al.* [30] produces $\chi^2/\text{datum} = 37$ which is grossly unacceptable. In fact, only for the interval 0-8 MeV is NLO acceptable. This range of validity is so tiny that no serious implications can be drawn from any prediction by this potential.

Finally, we like to take this opportunity to also present an overview of other interesting Nd observables, which are shown in Figs. 2-8. Concerning the deuteron vector analyzing power, iT_{11}, at 3 and 10 MeV (Fig. 2), it should be noted that a seemingly drastic reduction

of the discrepancy between theory and pd data seen at 3 MeV has its origin in large effects of the long range Coulomb force acting between two protons, which is not taken into account in our calculations [6]. Taking this Coulomb force into account, iT_{11} is also underpredicted in the peak region [6]—an equally well-known problem, which is why it would be appropriate to speak more generally of the vector analyzing power puzzle in elastic Nd scattering. In almost all cases, the Idaho chiral NN potential follows the trend of the high-precision potentials. The only exceptions are T_{20} and T_{21} at 65 MeV where the chiral potential predictions describe slightly better the data in the minimum region as compared to conventional potentials. But, apart from this, the quantitative chiral NN model containing contributions of higher orders in power counting does not produce any new signatures.

In summary, our main conclusions are:

- A *quantitative* chiral two-nucleon potential does not resolve the Nd A_y puzzle on the two-body force level.
- Low-energy ${}^3P_J$ NN phase shifts that "solve" the Nd A_y puzzle are inconsistent with the low-energy NN data.

And, finally, as a consequence of the above two points, one may expect that *no quantitative two-nucleon force—no matter what the basis is, pure phenomenology, meson theory, chiral EFT, or anything—will ever solve the Nd A_y puzzle.*

An accurate NN model requires to take chiral perturbation theory (χPT) to order four. At that order, also many three-body force (3NF) terms occur. According to the basic rules of χPT, all two- and many-body terms must be included for a complete calculation. Conventional 3NFs were shown to be ineffective for the A_y problem [14,6]. The advantage of χPT is that it provides a systematic scheme to generate all terms at a given order. As it turns out, there are several such chiral 3NF terms that were never considered in few-nucleon physics before [15–18]. It is natural to expect the resolution of the Nd A_y puzzle from such new chiral 3NFs which, therefore, should be at the focus of future work in the field.

At next-to-leading order (NLO) in χPT, there are no 3NF contributions. So, a calculation with a NLO two-nucleon potential and no 3NF seems to have formal validity. However, since at NLO the NN data can only be reproduced for $T_{\rm lab} \leq 8$ MeV, such a calculation is doomed to be inconclusive, from the outset, and higher order terms must be taken into account for any meaningful calculation.

ACKNOWLEDGMENTS

This work was supported in part by the U.S. National Science Foundation under Grant No. PHY-0099444, by the Ramón Areces Foundation (Spain), and by the U.S. Department of Energy, Office of High Energy and Nuclear Physics, under Grant No. DE-FG02-97ER41033. The numerical calculations have been performed on the CRAY T90 of the John von Neumann Institute for Computing in Jülich, Germany.

[1] C. Stolk and J. A. Tjon, Nucl. Phys. **A295**, 384 (1978).
[2] R. V. Reid, Ann. Phys. (N.Y.) **50**, 411 (1968).
[3] M. Lacombe, B. Loiseau, J. M. Richard, R. Vinh Mau, J. Côté, P. Pires, and R. de Tourreil, Phys. Rev. **21**, 861 (1980).
[4] W. Plessas and J. Haidenbauer, Few-Body Syst., Suppl. **2**, 185 (1987); Y. Koike and J. Haidenbauer, Nucl. Phys. **A463**, 365c (1987); J.Haidenbauer, Y. Koike, W. Plessas and H. Zankel, Proc. Intern. Workshop on Few-Body Approaches to Nuclear Reactions in Tandem and Cyclotron Energy Regions, edited by S. Oryu and T. Sawada (World Scientific, Singapore, 1987), p. 23.
[5] H. Witała, W. Glöckle, and T. Cornelius, Few-Body Syst., Suppl. **2**, 555 (1987); Nucl. Phys. **A491**, 157 (1989).
[6] A. Kievsky, M. Viviani, and S. Rosati, Nucl. Phys. **A577**, 511 (1994); Phys. Rev. C **52**, R15 (1995).
[7] W. Glöckle, H. Witała, D. Hüber, H. Kamada, and J. Golak, Phys. Rep. **274**, 107 (1996).
[8] S. C. Pieper, Nucl. Phys. **A193**, 529 (1972).
[9] P. Doleschall, Phys. Lett. **38B**, 298 (1972); Nucl. Phys. **A201**, 264 (1973); *ibid.* **A220**, 491 (1974); P. Doleschall, W. Grüebler, V. König, P. A. Schmelzbach, F. Sperisen, and B. Jenny, Nucl. Phys. **A380**, 72 (1982).
[10] H. Witała, W. Glöckle, and T. Cornelius, Nucl. Phys. **A496**, 446 (1989).
[11] H. Witala and W. Glöckle, Nucl. Phys. **A528**, 48 (1991).
[12] W. Tornow, H. Witała, and A. Kievsky, Phys. Rev. C **57**, 555 (1998); W. Tornow, and H. Witała, Nucl. Phys. **A637**, 280 (1998); T. Tornow and W. Tornow, Few-Body Syst. **26**, 1 (1999).
[13] D. Hüber and J. L. Friar, Phys. Rev. C **58**, 674 (1998).
[14] D. Hüber, H. Witała, and W. Glöckle, Few-Body Syst. **14**, 171 (1993); H. Witała, D. Hüber, and W. Glöckle, Phys. Rev. C **49**, R14 (1994).
[15] D. Hüber, J. L. Friar, and U. van Kolck, Phys. Rev. C **59**, 53 (1999).
[16] D. Hüber, J. L. Friar, A. Nogga, H. Witała, and U. van Kolck, Few-Body Syst. **30**, 95 (2001).
[17] J. L. Friar, Nucl. Phys. **A684**, 200 (2001).
[18] L. Canton and W. Schadow, Phys. Rev. C **64**, 031001 (2001).
[19] F. Sammarruca, H. Witała, and X. Meng, Acta Physica Polonica B **31**, 2039 (2000).
[20] E. Epelbaum, H. Kamada, A. Nogga, H. Witała, W. Glöckle, and U.-G. Meißner, Phys. Rev. Lett. **86**, 4787 (2001).
[21] S. Weinberg, Phys. Lett. B **251**, 288 (1990); Nucl. Phys. **B363**, 3 (1991).
[22] C. Ordóñez and U. van Kolck, Phys. Lett. B **291**, 459 (1992).
[23] C. Ordóñez, L. Ray, and U. van Kolck, Phys. Rev. Lett. **72**, 1982 (1994); Phys. Rev. C **53**, 2086 (1996).
[24] U. van Kolck, Prog. Part. Nucl. Phys. **43**, 337 (1999).
[25] L. S. Celenza, A. Pantziris, and C. M. Shakin, Phys. Rev. C **46**, 2213 (1992).
[26] C. A. da Rocha and M. R. Robilotta, Phys. Rev. C **49**, 1818 (1994); *ibid.* **52**, 531 (1995); M. R. Robilotta, Nucl. Phys. **A595**, 171 (1995); M. R. Robilotta, and C. A. da Rocha, Nucl. Phys. **A615**, 391 (1997); J.-L. Ballot, C. A. da Rocha, and M. R. Robilotta, Phys. Rev. C **57**, 1574 (1998).
[27] N. Kaiser, R. Brockmann, and W. Weise, Nucl. Phys. **A625**, 758 (1997).
[28] N. Kaiser, S. Gerstendörfer, and W. Weise, Nucl. Phys. **A637**, 395 (1998).

[29] N. Kaiser, Phys. Rev. C **61**, 014003 (1999); *ibid.* **62**, 024001 (2000); *ibid.* **63**, 044010 (2001); nucl-th/0107064.
[30] E. Epelbaum, W. Glöckle, and U.-G. Meißner, Nucl. Phys. **A637**, 107 (1998); *ibid.* **A671**, 295 (2000).
[31] D. B. Kaplan, M.J. Savage, and M.B. Wise, Nucl. Phys. **B478**, 629 (1996); D. B. Kaplan, Nucl. Phys. **B494**, 471 (1997); D. B. Kaplan, M.J. Savage, and M.B. Wise, Nucl. Phys. **B534**, 329 (1998); Phys. Lett. **B424**, 390 (1998).
[32] R. J. Furnstahl, B. D. Serot, and H.-B. Tang, Nucl. Phys. **A615**, 441 (1997); J. V. Steel and R. J. Furnstahl, *ibid.* **A637**, 46 (1998); R. J. Furnstahl, J. V. Steel, and N. Tirfessa, *ibid.* **A671**, 396 (2000); R. J. Furnstahl, H. W. Hammer, and N. Tirfessa, *ibid.* **A689**, 846 (2001).
[33] T.-S. Park, K. Kubodera, D. P. Min, and M. Rho, Phys. Rev. C **58**, 637 (1998); Nucl. Phys. **A646**, 83 (1999).
[34] T. D. Cohen, Phys. Rev. C **55**, 67 (1997); D. R. Phillips and T. D. Cohen, Phys. Lett. **B390**, 7 (1997); K. A. Scaldeferri, D. R. Phillips, C. W. Kao, and T. D. Cohen, Phys. Rev. C **56**, 679 (1997); S. R. Beane, T. D. Cohen, and D. R. Phillips, Nucl. Phys. **A632**, 445 (1998).
[35] G. Rupak and N. Shoresh, nucl-th/9906077; P. F. Bedaque, H. W. Hammer, and U. van Kolck, Nucl. Phys. **A676**, 357 (2000); S. Fleming, Th. Mehen, and I. W. Stewart, Nucl. Phys. **A677**, 313 (2000); Phys. Rev. C **61**, 044005 (2000).
[36] S. R. Bean, P. F. Bedaque, M. J. Savage, and U. van Kolck, nucl-th/0104030.
[37] V. G. J. Stoks, R. A. M. Klomp, M. C. M. Rentmeester, and J. J. de Swart, Phys. Rev. C **48**, 792 (1993).
[38] R. Machleidt, F. Sammarruca, and Y. Song, Phys. Rev. C **53**, 1483 (1996).
[39] R. Machleidt, Phys. Rev. C **63**, 024001 (2001).
[40] D. R. Entem and R. Machleidt, nucl-th/0107057; nucl-th/0108057, Phys. Lett. B, in press.
[41] V. G. J. Stoks, R. A. M. Klomp, C. P. F. Terheggen, and J. J. de Swart, Phys. Rev. C **49**, 2950 (1994).
[42] R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, Phys. Rev. C **51**, 38 (1995).
[43] The 1999 world *np* database is the Nijmegen 1992 *np* database [37] plus all *np* data published between January 1993 and December 1999 and not rejected in the analysis. The latter data are listed in Table XVI of Ref. [39].
[44] J. E. McAninch, W. Haeberli, H. Witała, W. Glöckle, J. Golak, Phys. Lett. B **153**, 29 (1985).
[45] S. Shimizu, K. Sagara, H. Nakamura, K. Maeda, T. Miwa, N. Nishimori, S. Ueno, T. Nakashima, and S. Morinobu, Phys. Rev. C **52**, 1193 (1995).
[46] W. Tornow, C. R. Howell, R. C. Byrd, R. S. Pedroni, R. L. Walter, Phys. Rev. Lett. **49**, 312 (1982).
[47] F. Sperisen, W. Grüebler, V. Koenig, P. A. Schmelzbach, K. Elsener, B. Jenny, C. Schweizer, J. Ulbricht, P. Doleschall, Nucl. Phys. A **422**, 81 (1984).
[48] H. Rühl, B. Dechant, J. Krug, W. Lübcke, G. Spangardt, M. Steinke, M. Stephan, D. Kamke, J. Balewski, K. Bodek, L. Jarczyk, A. Strzałkowski, W. Hajdas, St. Kistryn, R. Müller, J. Lang, R. Henneck, H. Witała, Th. Cornelius, W. Glöckle, Nucl. Phys. A **524**, 377 (1991).
[49] H. Shimizu, K. Imai, N. Tamura, K. Nisimura, K. Hatanaka, T. Saito, Y. Koike, Y. Taniguchi, Nucl. Phys. A **382**, 242 (1982).
[50] K. Sagara, H. Ogari, S. Shimizu, K. Maeda, H. Nakamura, T. Nakashima, S. Morinobu, Phys. Rev. C **50**, 576 (1994).
[51] H. Witała, W. Glöckle, L. E. Antonuk, J. Arvieux, D. Bachelier, B. Bonin, A. Boudard, J. M.

Cameron, H. W. Fielding, M. Garcon, F. Jourdan, C. Lapointe, W. J. McDonald, J. Pasos, G. Roy, I. The, J. Tinslay, W. Tornow, J. Yonnet, W. Ziegler, Few-Body Systems **15**, 67 (1993).

[52] G. Rauprich, S. Lemaitre, P. Niessen, K. R. Nyga, R. Reckenfelderbäumer, L. Sydow, H. Paetz gen. Schieck, H. Witała, W. Glöckle, Nucl. Phys. A **535**, 313 (1991).

[53] H. R. Setze, C. R. Howell, W. Tornow, R. T. Braun, W. Glöckle, A. H. Hussein, J. M. Lambert, G. Mertens, C. D. Roper, F. Salinas, I. Slaus, D. E. González Trotter, B. Vlahović, R. L. Walter, H. Witała, Phys. Lett. B **388**, 229 (1996).

[54] J. Strate, K. Geissdörfer, R. Lin, J. Cub, E. Finckh, K. Gebhardt, S. Schindler, H. Witała, W. Glöckle, Th. Cornelius, J. Phys. G: Nucl. Phys. **14**, L 229 (1988).

TABLE I. $^1S_0$ np and $^3S_1$ phase shifts (in degrees).

$T_{\rm lab}$ (MeV)	PWA93[a]	NLO[b]	Idaho[c]	CD-Bonn[d]
		$^1\boldsymbol{S_0}$		
1	62.068(30)	62.05	62.03	62.09
5	63.63(8)	63.85	63.52	63.67
10	59.96(11)	60.26	59.80	60.01
25	50.90(19)	50.92	50.66	50.93
50	40.54(28)	39.29	40.27	40.45
100	26.78(38)	21.68	26.87	26.38
150	16.94(41)	7.26	17.56	16.32
200	8.94(39)	-5.53	9.58	8.31
		$^3\boldsymbol{S_1}$		
1	147.747(10)	147.70	147.76	147.75
5	118.178(21)	118.29	118.20	118.18
10	102.611(35)	102.84	102.64	102.62
25	80.63(7)	80.69	80.67	80.63
50	62.77(10)	61.74	62.80	62.73
100	43.23(14)	38.65	43.13	43.06
150	30.72(14)	21.67	30.41	30.47
200	21.22(15)	7.20	20.90	20.95

[a]Nijmegen multi-energy np analysis [37]. Numbers in parentheses give the uncertainties in the last digits.
[b]Chiral next-to-leading order (NLO) potential by Epelbaum *et al.* [30] using a Gaussian cutoff with cutoff mass $\Lambda = 600$ MeV.
[c]Chiral NN potential by Entem and Machleidt [40] ('Idaho-B' is used).
[d]Reference [39].

TABLE II. Triplet-P np phase shifts (in degrees). For notation see Table I.

$T_{\rm lab}$ (MeV)	PWA93[a]	NLO	Idaho	Modified[b]	CD-Bonn
			$^3P_0$		
1	0.18	0.19	0.18	0.18	0.18
5	1.63(1)	1.67	1.64	1.61	1.61
10	3.65(2)	3.72	3.69	3.62	3.62
25	8.13(5)	8.22	8.19	7.98	8.10
50	10.70(9)	10.84	10.63	10.28	10.74
100	8.46(11)	8.31	8.17	7.91	8.57
150	3.69(14)	2.52	3.60	3.66	3.72
200	-1.44(17)	-4.10	-1.21	-0.93	-1.55
			$^3P_1$		
1	-0.11	-0.12	-0.11	-0.11	-0.11
5	-0.94	-0.99	-0.93	-0.94	-0.93
10	-2.06	-2.16	-2.04	-2.09	-2.04
25	-4.88(1)	-5.03	-4.82	-4.98	-4.81
50	-8.25(2)	-8.32	-8.22	-8.57	-8.18
100	-13.24(3)	-12.66	-13.36	-13.86	-13.23
150	-17.46(5)	-15.94	-17.59	-17.85	-17.51
200	-21.30(7)	-18.86	-21.28	-21.18	-21.38
			$^3P_2$		
1	0.02	0.02	0.02	0.02	0.02
5	0.25	0.24	0.26	0.28	0.26
10	0.71	0.70	0.72	0.78	0.72
25	2.56(1)	2.89	2.58	2.78	2.60
50	5.89(2)	8.29	5.86	6.29	5.93
100	10.94(3)	22.61	10.77	11.24	11.01
150	13.84(4)	35.98	13.72	13.82	13.98
200	15.46(5)	44.31	15.58	15.33	15.66

[a]Uncertainties smaller than 0.005 degrees are not shown.
[b]Modified version of the Idaho potential with enhanced spin-orbit force at low energies.

TABLE III. Spin-orbit phase shift combination $\Delta_{\rm LS}$ (in degrees), Eq. (1), at 10, 25, and 50 MeV for various models explained in Table I and II.

$T_{\rm lab}$ (MeV)	PWA93	NLO	Idaho	Modified	CD-Bonn
10	0.203	0.212	0.195	0.244	0.207
25	0.93	1.09	0.92	1.07	0.94
50	2.73	3.73	2.73	3.05	2.73

TABLE IV. χ^2/datum for the reproduction of the 1999 *np* database [43] up to $T_{\rm lab} = 210$ MeV by various models explained in Table I and II.

Bin (MeV)	# of data	PWA93	NLO	Idaho	Modified	CD-Bonn
0-8	81	1.05	1.05	1.03	1.32	1.06
8-17	192	0.86	1.22	0.85	1.87	0.91
17-35	292	0.82	2.15	0.81	1.27	0.81
35-75	340	1.03	10.8	1.09	1.24	1.01
75-125	239	1.00	11.6	0.99	1.06	0.97
125-183	414	1.06	95.7	1.07	1.15	1.03
183-210	141	0.96	111.7	0.97	1.18	1.01
0-210	1699	0.97	37.0	0.98	1.27	0.97



FIG. 1. Nucleon analyzing power A_y for elastic Nd scattering at 3, 10, and 65 MeV. The shaded band represents the predictions by the high-precision potentials CD-Bonn [39], Argonne V_{18} [42], and the Nijmegen potentials Nijm-I, Nijm-II, and Nijm93 [41] (using always the np versions of these potentials). The dashed line is the prediction by the Idaho chiral NN potential [40]. The dotted curve represents the result from the 'Modified' chiral potential (see text) and the dash-dot line is predicted by the NLO chiral potential by Epelbaum *et al.* [30]. Data at 3 MeV are from [44] (nd, squares), at 10 MeV from [46] (nd, squares), and at 65 MeV from [48] (nd, squares) and [49] (pd, crosses).



FIG. 2. Deuteron vector analyzing power iT_{11} for elastic Nd scattering. Energies, bands, and curves as in Fig. 1. The circles are pd data: at 3 MeV from [45], at 10 MeV from [47], and at 65 MeV from [51].



FIG. 3. Tensor analyzing power T_{20} for elastic Nd scattering. Energies, bands, and curves as in Fig. 1. The circles are pd data: at 3 MeV from [45], at 10 MeV from [47], and at 65 MeV from [51].



FIG. 4. Tensor analyzing power T_{21} for elastic Nd scattering. Energies, bands, and curves as in Fig. 1. The circles are pd data: at 3 MeV from [45], at 10 MeV from [47], and at 65 MeV from [51].



FIG. 5. Tensor analyzing power T_{22} for elastic Nd scattering. Energies, bands, and curves as in Fig. 1. The circles are pd data: at 3 MeV from [45], at 10 MeV from [47], and at 65 MeV from [51].



FIG. 6. Spin transfer coefficients $K_x^{x'}$, $K_y^{y'}$, and $K_z^{x'}$ for elastic Nd scattering at 10 MeV. Bands and curves as in Fig. 1. The circles are pd data from [47].



FIG. 7. Differential cross section $d\sigma/d\Omega$ of elastic Nd scattering. Energies, bands, and curves as in Fig. 1. The crosses are pd data: at 3 MeV from [50], at 10 MeV from [47], and at 65 MeV from [49]. The circles at 65 MeV are nd data from [48].



FIG. 8. Neutron-deuteron breakup cross sections for the space-star and quasi-free scattering (QFS) configurations at 13 MeV along the kinematical locus S. Bands and curves as in Fig. 1. The circles are *pd* data from [52]. The crosses and squares are *nd* data from [53] and [54], respectively.